arXiv:1702.05816v1 [math.DG] 19 Feb 2017

THE ISOPERIMETRIC PROBLEM FOR LENS SPACES

CELSO VIANA

Abstract. We solve the isoperimetric problem in the Lens spaces with large fundamental group. Namely, we prove that the isoperimetric surfaces are geodesic spheres or tori of revolution about geodesics. We also show that the isoperimetric problem in $L(3,1)$ and $L(3,2)$ follows from the proof of the Willmore conjecture by Marques and Neves.

1. Introduction

The isoperimetric problem is a classical subject in Differential Geometry with its origin in ancient Greece. It consists in finding on a Riemannian manifold M the regions that minimize the perimeter among sets enclosing the same volume. The solutions are called isoperimetric regions and their boundaries isoperimetric hypersurfaces. The Euclidean plane is historically the first space where the problem started to be investigated rigorously. It is now a well known fact that the round circles are the optimal curves for the problem. This geometric fact is often seen through the following classical inequality:

$$L^2 \geq 4\pi A, \tag{1.1}$$

where L and A stand for the length and the enclosed area of a simple closed curve $\gamma : \mathbb{S}^1 \to \mathbb{R}^2$ respectively.

The framework of geometric measure theory and its tools work successfully well in tackling the aspects of existence and regularity of this variational problem. When M^{n+1} is closed or homogeneous, then isoperimetric hypersurfaces do exist and are smooth up to a closed set of Hausdorff dimension $n-7$. The regular part is a stable hypersurface of constant mean curvature. This major contribution to the isoperimetric problem was achieved thanks to the efforts of many people, including F. Almgren, R. Schoen, L. Simon, F. Morgan, and others (see [M] for a comprehensive list). Despite the long history of the problem, it remains largely open with few 3-manifolds where the problem is completely understood.

C.V. was supported by The EPSRC Centre for Doctoral Training in Geometry and Number Theory: The London School of Geometry and Number Theory.

The simply connected space forms, $\mathbb{S}^{n+1}$, $\mathbb{R}^{n+1}$ and $\mathbb{H}^{n+1}$, are the most appealing spaces to begin the study of the isoperimetric problem. It turns out that their symmetries are enough to characterize the geodesic spheres as the isoperimetric hypersurfaces.

A complete solution on $\mathbb{S}^2 \times \mathbb{S}^1$ with the standard product metric can be found in [PR]. For other homogeneous manifolds with certain product structure, such as $\mathbb{H}^2 \times \mathbb{R}$, $\mathbb{H}^2 \times \mathbb{S}^1$, and $\mathbb{S}^n \times \mathbb{R}$, see [HH], [PR], and [P] respectively. The case $\mathbb{S}^1 \times \mathbb{R}^n$ is also treated in [PR] and they show that when $n \geq 9$ unduloids are minimizers rather than cylinders for certain volumes. One key idea exploited in the results listed above is the use of symmetry to reduce the problem to an ODE analysis. The case $T^2 \times \mathbb{R}$ where T^2 is a flat torus is not solved in full; great progress can be found in [RR1] and in [HPRR]. More generally, it is known that boundaries of small isoperimetric regions in closed manifolds are nearly round spheres, see [MJ]. To finish this brief and not exhaustive account of results on the isoperimetric problem we mention that Bray and Morgan ([B] and [BM]) classified the horizon homologous isoperimetric surfaces in the Schwarzschild manifold. The work in [B] highlighted the interesting relationship between isoperimetric surfaces and the concept of mass in general relativity.

We will be interested in spherical space forms in this paper. A significant result in this direction was given in [RR] where A. Ros and M. Ritoré solved the isoperimetric problem in the projective space $\mathbb{RP}^3$. They show that the solutions are geodesic spheres or flat tori. Later, A. Ros [R] used the above result to give a proof of the Willmore conjecture in $\mathbb{S}^3$ for the special case of surfaces that are invariant by the antipodal map.

The real projective space is a special case of an important family of Riemannian manifolds, namely the Lens spaces $L(p, q)$. These are elliptic space forms obtained as a quotient of $\mathbb{S}^3$ by a finite group of isometries that are isomorphic to $\mathbb{Z}_p$ but which also depend on q. They are, along with $\mathbb{S}^2 \times \mathbb{S}^1$, characterized by having Heegaard genus one.

We give a complete solution for the isoperimetric problem in the Lens spaces with large fundamental group:

Theorem 1.1. *There exists a positive integer p_0 such that for every $p \geq p_0$ and every $q \geq 1$ the isoperimetric surfaces in $L(p, q)$ are either geodesic spheres or tori of revolution about geodesics.*

The isoperimetric problem in dimension three was previously solved for only a finite number of non-diffeomorphic 3-manifolds.

We also add to the literature the observation that the proof of the Willmore conjecture by Marques and Neves [MN] can be applied to

extend the work of Ros and Ritoré [RR], on the classification of stable CMC surfaces in $\mathbb{RP}^3$, to $L(3,1)$ and $L(3,2)$: [1]

Theorem 1.2. *The immersed stable CMC surfaces in $L(3,1)$ and $L(3,2)$ are either geodesic spheres or flat tori. Moreover, the minimal Clifford torus is, up to ambient isometries, the only index one minimal surface in $L(3,1)$ and $L(3,2)$.*

The idea of the proof of Theorem 1.1 is as follows. Stability implies that every isoperimetric surface is connected and its genus is $0, 1, 2$ or 3. If follows from a classical result of Hopf that if the genus is 0, then it is a geodesic sphere. From [RR] we know that if the genus is 1, then it is flat, and this forces the surface to be of revolution about a geodesic. We are left to rule out other topological types. To do so we argue by contradiction. We assume that there exists a sequence of Lens spaces with increasing fundamental group containing isoperimetric surfaces of genus 2 or 3. After a suitable rescaling on the metrics we use compactness results to obtain a limit for the sequence of Lens spaces which will be a flat three manifold of rank one. In the same way, the sequence of isoperimetric surfaces will converge to a flat surface in the respective ambient manifold. The topology of the surfaces will force the limit to be an union of planes. On the other hand, the minimization property rules this configuration out.

The arguments in the proof of Theorem 1.1 generalize naturally for the Berger Spheres $\mathbb{S}^3_\varepsilon$. This is a well known one parameter family of homogeneous metrics on the 3-sphere; the case $\varepsilon = 1$ corresponds to the round metric. All the Hopf fibers in $\mathbb{S}^3_\varepsilon$ have the same length $2\pi\,\varepsilon$. When ε is not too small, spheres are the only solutions of the isoperimetric problem, see [TU]. When ε is very small, it is pointed out in [TU] that some tori are better candidates to solve the isoperimetric problem rather than spheres for certain volumes.

Theorem 1.3. *There exists $\varepsilon_0 > 0$ such that for every $\varepsilon < \varepsilon_0$ the isoperimetric surfaces in the Berger spheres $\mathbb{S}^3_\varepsilon$ are either rotationally invariant spheres or tori.*

Remark 1.4. It follows from the work in [MHH] and [PR] that the minimal Clifford torus in $L(p,1)$ (resp. $\mathbb{S}^3_\varepsilon$) is isoperimetric for every $p \geq 3$ (resp. $\varepsilon \leq \frac{1}{3}$), see Section 3.

Acknowledgements. I am grateful to my advisor André Neves for his constant encouragement and support during the course of this work. I also thank Jason Lotay for many helpful comments on an earlier version of this manuscript.

[1] $L(3,1)$ and $L(3,2)$ are diffeomorphic but not isometric.

2. Preliminaries

Let (M^{n+1}, g) be an orientable Riemannian manifold of dimension $n+1$. The $n+1$-dimensional Hausdorff measure of a region $\Omega \subset M$ is denoted by $|\Omega|$. Similarly, we denote the n-dimensional Hausdorff measure of the hypersurface $\partial\Omega \subset M$ by $|\partial\Omega|$. The class of regions considered here are those of finite perimeter, see [S].

2.1. **Isoperimetric.** A region $\Omega \subset M$ is called an *Isoperimetric region* if

$$|\partial\Omega| = \inf\{|\partial\Omega'| \,:\, \Omega' \subset M \quad \textit{and} \quad |\Omega'| = |\Omega|\}.$$

In this case, the hypersurface $\Sigma = \partial\Omega$ is called an *Isoperimetric hypersurface*.

In general, the existence of isoperimetric surfaces is handled by a compactness theorem from geometric measure theory. For non-compact manifolds one needs to be careful since a minimizing sequence of regions of fixed volume may drift off to infinity. We recommend [M] for a recent reference on the regularity of isoperimetric hypersurfaces:

Theorem 2.1. *Let (M^{n+1}, g) be a closed Riemannian manifold. For every $0 < t < vol(M)$ there exists an isoperimetric region Ω satisfying $|\Omega| = t$. Moreover, $\Sigma = \partial\Omega$ is smooth up to a closed set of Hausdorff dimension $n-7$.*

2.2. **Stability.** Isoperimetric hypersurfaces are stable critical points of the area functional for variations that preserve the enclosed volume; thus, the regular part of isoperimetric hypersurfaces has constant mean curvature. More generally, we say an isometric immersion $\phi : \Sigma^n \to M^{n+1}$ has *constant mean curvature* (CMC) if it is a critical point of the area functional for volume preserving variations. Critical points are called *stable CMC* if the second derivative of the area is non-negative for such variations.

Equivalently, ϕ is stable if for every $f \in C^\infty(\Sigma)$ with compact support such that $\int_\Sigma f \, dvol_\Sigma = 0$, we have

(2.1)

$$I(f,f) = -\int_\Sigma f L f \, d_\Sigma := \int_\Sigma |\nabla f|^2 - (Ric(N,N) + |A|^2) f^2 \, d_\Sigma \geq 0.$$

N is the unit normal vector of Σ and A is the *second fundamental form* of the immersion ϕ. The *mean curvature* of Σ, denoted by H, is defined by $2H = trace(A)$.

The study of immersed stable CMC hypersurfaces started in [BC] and [BCE] with a new characterization of the geodesic spheres in the simply connected space forms $\mathbb{R}^{n+1}$, $\mathbb{S}^{n+1}$ and $\mathbb{H}^{n+1}$. The classification

of stable CMC surfaces is often a way to approach the isoperimetric problem in reasonable spaces. With this purpose in mind, A. Ros and M. Ritoré in [RR] used the Hersch-Yau trick to study orientable stable CMC surfaces on three manifolds of positive Ricci curvature.

Theorem 2.2 (Ros-Ritoré [RR])**.** *Let (M, g) be a closed three manifold with positive Ricci curvature. If $\phi : \Sigma \to M^3$ is a stable CMC immersion, then $g(\Sigma) \leq 3$. Moreover, if $g(\Sigma) = 2$ or 3, then*

$$\big(\frac{1}{2}\inf_{\Sigma} Ric_M + H^2\big)|\Sigma| \leq 2\pi.$$

Proposition 2.3. *Let $\phi : \Sigma \to M^3$ be a stable immersion with constant mean curvature H into an elliptic space form $M = \mathbb{S}^3/G$. Then*

(1) If $g(\Sigma) = 2$ or 3, then $\big(1 + H^2\big)|\Sigma| \leq 2\pi$.
(2) If $g(\Sigma) = 2$ or 3 and $|G| \leq 4$, then ϕ is an embedding. Moreover, if $|G| \leq 6$, then the pullback of Σ, through the covering map $\Pi : \mathbb{S}^3 \to M^3$, is connected.
(3) If $|G| = 2$ or 3, then $g(\Sigma) = 0$ or 1.

Proof. The first statement follows from the previous theorem since $Ric_M = 2$. Let $\phi_* : \pi_1(\Sigma) \to \pi_1(M)$ be the induced map in fundamental groups. As $K = Ker(\phi_*)$ has finite index there exists a finite covering $\psi : \widetilde{\Sigma} \to \Sigma$ such that $Im\big(\psi_*\big) = K$ and $\big(\phi \circ \psi\big)_* = 0$. This means there exists a lifting of this map into $\mathbb{S}^3$ and we denote it by $\widetilde{\phi} : \widetilde{\Sigma} \to \mathbb{S}^3$. It follows that $\big(1 + H^2\big)|\widetilde{\Sigma}| \leq |G|\, 2\pi$. If ϕ is not an embedding, then $\widetilde{\phi}$ is not embedding either. By the work of Li and Yau [LY] the Willmore energy of the immersed surface $\widetilde{\Sigma}$, i.e. $\mathcal{W}(\widetilde{\Sigma}) = \int_{\widetilde{\Sigma}} \big(1 + H^2\big) dvol_{\Sigma}$, is strictly greater[2] than 8π. Therefore, if $|G| \leq 4$, we obtain a contradiction and $\widetilde{\phi}$ is an embedding. Moreover, for closed surfaces with genus greater than or equal to 1 in $\mathbb{S}^3$ the Willmore conjecture, recently proved in [MN], states that $\mathcal{W}(\Sigma) \geq 2\pi^2$. Let $\cup_{i=1}^{l} \widetilde{\Sigma}_i$ be the pre-image of Σ by the universal covering map, then

$$2\, l\, \pi^2 \leq \sum_{i=1}^{l} \mathcal{W}\big(\widetilde{\Sigma}_i\big) = |G|\mathcal{W}(\Sigma) \leq |G| 2\pi \Rightarrow \frac{|G|}{l} \geq \pi$$

Therefore, if $|G| \leq 6$, then $l = 1$ and $|G| \geq \pi$. In particular, if $|G| = 2$ or 3, then there exist no stable CMC surface Σ with $g(\Sigma) \geq 2$. □

Definition 2.2. For each $r \in (0, \frac{\pi}{2})$ we define the Clifford Torus T_r as:

$$T_r = \mathbb{S}^1(\cos(r)) \times \mathbb{S}^1(\sin(r)) \subset \mathbb{S}^3. \tag{2.3}$$

[2]The case $\mathcal{W}(\Sigma) = 8\pi$ is discussed in [RR].

Every flat tori with constant mean curvature in $\mathbb{S}^3$ is congruent to a Clifford torus T_r. This follows from the Rigidity theorem, pg. 49 in [C], for they have the same second fundamental form.

Corollary 2.4. *The stable CMC surfaces in $L(3,1)$ and $L(3,2)$ are totally umbilical spheres or flat tori. In addition, the index one minimal surfaces in $L(3,1)$ and $L(3,2)$ are congruent to the the projection of minimal Clifford torus.*

Proof. Let $\Sigma \subset L(3,q)$, $q = 1, 2$, be in the conditions of the corollary. By Proposition 2.3, $g(\Sigma) = 0$ or 1. If $g(\Sigma) = 0$, then it follows from the Hopf holomorphic quadratic differential that Σ is totally umbilical. If $g(\Sigma) = 1$, then it is proved in [RR] that Σ is flat. □

2.3. **Isoperimetric profile.** The isoperimetric properties of M can be encapsulated in a single function called the *Isoperimetric profile*. This is the function $I_M : [0, \mathrm{vol}(M)] \to [0, +\infty)$ defined by

$$I_M(v) = \inf\{|\partial\Omega| \,:\, \Omega \subset M \quad \text{and} \quad |\Omega| = v\}. \tag{2.4}$$

We finish the section with some well known facts on the analytic nature of I_M. These will be used later in Section 3.

Let Ω be an isoperimetric region in M such that $|\Omega| = v$ for some $v \in (0, \mathrm{Vol}(M))$. The function I_M has left and right derivative $(I_M)'_-(v)$ and $(I_M)'_+(v)$. In addition, if H is the mean curvature of $\Sigma = \partial\Omega$ in the direction of the inward unit vector, then

$$(I_M)'_+(v) \leq 2H \leq (I_M)'_-(v). \tag{2.5}$$

The second derivative also exists but weakly in the sense of comparison functions. More precisely, we say $f'' \leq h$ weakly at x_0 if there exists a smooth function g such that $f \leq g$, $f(x_0) = g(x_0)$, and $g'' \leq h$. In this sense we have

$$I_M(v)^2\, I''_M(v) + \int_\Sigma \left(Ric_g(N,N) + |A|^2\right) d_\Sigma \leq 0. \tag{2.6}$$

The equations (2.5) and (2.6) are first presented on [BP]. We sketch the proof of (2.5) and (2.6).

Let Σ_V be the variation $\Sigma_t = exp_\Sigma(tN)$ of Σ reparametrized in terms of the enclosed volume $v(t)$. In addition, let $\phi(t)$ (resp. $\phi(v)$) be the area of Σ_t (resp. Σ_v). By the first variation formula for the area and volume we have $\phi'(0) = 2\,H\,|\Sigma|$ and $v'(0) = |\Sigma|$ respectively. Since $\phi'(t) = \phi'(v)v'(t)$, we conclude that $\phi'(v(0)) = 2H$ and also that $v'(0)^2\phi''(v(0)) = \phi''(0) - \phi'(v(0))v''(0)$. On the other hand, the

second derivative of area for general variations implies the following:

$$\begin{aligned}\phi''(0) &= -\int_\Sigma 1\, L\, 1\, d_\Sigma + 2H\, v''(0)\\ &= -\int_\Sigma \big(Ric_g(N,N) + |A|^2\big) d_\Sigma + 2H\, v''(0).\end{aligned}$$

Hence, in the sense of comparison functions, (2.6) follows from:

$$\phi(v(0))^2\, \phi''(v(0)) + \int_\Sigma \big(Ric_g(N,N) + |A|^2\big) d_\Sigma = 0. \tag{2.7}$$

3. The Isoperimetric problem in the Lens Spaces

In order to define the Lens spaces, we first recall the round three sphere as:

$$\mathbb{S}^3 = \{(z,w) \in \mathbb{C}^2 \,:\, |z|^2 + |w|^2 = 1\}.$$

Fix p, q integers with the following property $1 \le q < p$ and $gcd(p,q) = 1$. Let $\mathbb{Z}_p$ be the group $\mathbb{Z}/p\mathbb{Z}$ acting on $\mathbb{S}^3$ as follows:

$$m \in \mathbb{Z}_p \mapsto m \cdot (z,w) = (e^{\frac{2\pi i q m}{p}} z, e^{\frac{2\pi i m}{p}} w). \tag{3.1}$$

The group $\mathbb{Z}_p$ acts freely and properly discontinuous on $\mathbb{S}^3$. The orbit space $\mathbb{S}^3/\mathbb{Z}_p$ is a closed three manifold called the *Lens space*, it is denoted by $L(p,q)$.

The Hopf fibration, which is the Riemannian submersion $h : \mathbb{S}^3 \to \mathbb{S}^2(\frac{1}{2})$ defined by $h(z,w) = \frac{z}{w}$, can be extended naturally to $L(p,q)$. Indeed, the group $\mathbb{Z}_p$ acts on the set of *Hopf fibers* through the following cyclic action on $\mathbb{S}^2(\frac{1}{2})$:

$$\Gamma_p = \langle e^{\frac{2\pi i(q-1)}{p}} \rangle : \lambda \in \mathbb{C} \cup \{\infty\} \mapsto e^{\frac{2\pi i(q-1)}{p}} \cdot \lambda \in \mathbb{C} \cup \{\infty\}.$$

The Hopf fibration for $L(p,q)$ is then defined as $h : L(p,q) \to \mathbb{S}^2(1/2)/\Gamma_p$. The set $\mathbb{S}^2(1/2)/\Gamma_p$ is a two dimensional orbifold with conical singularities at the north and south pole.

The preimage of horizontal slices of $\mathbb{S}^2(1/2)/\Gamma_p$ via h correspond to the Clifford torus described in (2.3). They are natural candidates to solve the Isoperimetric problem in $L(p,q)$.

3.1. **Comments on Steiner Symmetrization for Lens Spaces.**

Steiner and Schwarz symmetrization theorems were proved in [MHH] for certain fiber bundles such as the Lens spaces. To explain this symmetrization procedure we restrict to the case $L(p,1)$ where the Hopf fibration $h : L(p,1) \to \mathbb{S}^2(\frac{1}{2})$ is a smooth Riemannian submersion.

The symmetrization consist of associating to each set of finite perimeter $R \subset L(p,1)$ the set $Sym(R)$ in the product manifold $\mathbb{S}^2(\frac{1}{2}) \times \mathbb{S}^1(\frac{1}{p})$

defined by replacing the slice of R in each fiber with a ball of the same volume about the respective fiber in the product. The coarea formula for Riemannian submersions implies that $Sym(R)$ encloses the same amount of volume as R. It is proved in [MHH] that $Sym(R)$ has no greater perimeter than R.

One immediate consequence is that $I_{L(p,1)} \geq I_{\mathbb{S}^2(\frac{1}{2})\times\mathbb{S}^1(\frac{1}{p})}$. Applying the classification of the isoperimetric problem on $\mathbb{S}^2(\frac{1}{2})\times\mathbb{S}^1(\frac{1}{p})$, [PR], we conclude that $I_{L(p,1)} = I_{\mathbb{S}^2(\frac{1}{2})\times\mathbb{S}^1(\frac{1}{p})}$ in a interval around $V = \frac{\text{Vol}(L(p,1))}{2}$. In particular, the minimal Clifford torus is isoperimetric in $L(p,1)$ for every $p \geq 3$. The isoperimetric profiles, however, do not coincide as the profile of geodesic spheres on the respective spaces are different. Therefore, this technique is not enough to completely solve the isoperimetric problem.

It is important to point out that, for general Lens spaces $L(p,q)$, there is no analogue of [PR] for $\mathbb{S}^2(\frac{1}{2})/\Gamma_p \times \mathbb{S}^1(\frac{1}{n_p})$ which is a manifold having codimension two singularities.

3.2. **Some aspects of Lens Spaces.** For every $x \in L(p,q)$ the injectivity radius of $L(p,q)$ at x satisfies $\text{inj}_x L(p,q) \geq \frac{\pi}{p}$, with equality only at points in the critical fibers. Indeed, for $\theta = e^{\frac{2\pi i}{p}}$ we have:

$$\begin{aligned} a^2 := d^2_{\mathbb{R}^4}[(\theta^{kq}z, \theta^k w), (z,w)] &= |\theta^{kq}-1|^2|z|^2 + |\theta^k - 1|^2|w|^2 \geq |\theta - 1|^2 \\ d_{\mathbb{S}^3}[(\theta^{kq}z, \theta^k w), (z,w)] &= 2\arcsin(\frac{a}{2}) \geq 2\arcsin\big(\frac{2}{2}\sin\frac{2\pi}{2p}\big) = \frac{2\pi}{p}. \end{aligned}$$

However, is not true in general that $inj(L(p,q),x) = O(\frac{1}{p})$ as $p \to \infty$.

Example 3.1. Let's consider $L(k^2, k+1)$, $k \in \mathbb{Z}_+$. We show that the injectivity radius at points far away from the critical fibres are $O(\frac{1}{k})$. If the round metric is scaled by the factor k^2, then we have the Riemannian submersion:

$$h : \big(L(k^2,k+1), k^2 g_0, x_k\big) \to \big(\mathbb{S}^2/\mathbb{Z}_k, k^2 g_{\mathbb{S}^2}, h(x_k)\big).$$

The fibers have constant length 2π except the critical fibres which have length $\frac{2\pi}{p}$. The right hand side will converge, as $k \to \infty$, to $\mathbb{S}^1 \times \mathbb{R}$. It follows from the coarea formula for Riemannian submersions that the volume of the geodesic ball $B_{4\pi}(x_k)$ in $\big(L(k^2,k+1), k^2 g_0, x_k\big)$ is bounded from below. Therefore, by Cheeger's inequality, Lemma 51 in [PP], the injectivity radius of the sequence $\big(L(k^2,k+1), k^2 g_0, x_k\big)$ is bounded from below. This sequence converges to a flat $T^2 \times \mathbb{R}$.

If $x, y \in T_{\frac{\pi}{4}}/\mathbb{Z}_p \subset L(p,q)$, then $d_{L(p,q)}(x,y) \geq C d_{T_{\frac{\pi}{4}}/\mathbb{Z}_p}(x,y)$ for some constant $C > 0$ independent of p, q. Thus intrinsic and extrinsic distances on $T_{\frac{\pi}{4}}/\mathbb{Z}_p$ are equivalent.

Lemma 3.2. *If $x \in T_{\frac{\pi}{4}}/\mathbb{Z}_p \subset L(p,q)$ and the extrinsic diameter of $T_{\frac{\pi}{4}}/\mathbb{Z}_p$ in $L(p,q)$ is bounded from below, then $inj_x L(p,q) = O(\frac{1}{p})$.*

Proof. Let $\lambda_p = \frac{1}{inj_x L(p,q)}$ and recall that $\frac{1}{\lambda_p} \geq \frac{\pi}{p}$. Without loss of generality, let's assume that $\text{diameter}_{L(p,q)}(T_{\frac{\pi}{4}}/\mathbb{Z}_p) \geq 1$. Hence, under the rescaled metric $\lambda_p^2 g_{\mathbb{S}^3}$, the extrinsic diameter of $T_{\frac{\pi}{4}}/\mathbb{Z}_p$ is greater than or equal to λ_p. Let $\gamma_p(t)$ be a geodesic segment realizing the intrinsic diameter of $T_{\frac{\pi}{4}}/\mathbb{Z}_p$. Thus, we can find disjoint balls $B_R(x_i) \subset L(p,q)$, with $R < \frac{C}{2}$, $x_i \in \gamma_p(t)$, and $i = 1, \ldots, [\lambda_p] + 1$. Hence,

$$\sum_{i=1}^{[\lambda_p]+1} \mathcal{H}^2(B_R(x_{i_0}) \cap T_{\frac{\pi}{4}}/\mathbb{Z}_p) \leq |T_{\frac{\pi}{4}}/\mathbb{Z}_p| = \lambda_p^2 \frac{2\pi^2}{p}.$$

Therefore, there exists $i_0 \in \{1, \ldots, [\lambda_p] + 1\}$ such that:

$$C_1 \leq \mathcal{H}^2(B_R(x_{i_0}) \cap T_{\frac{\pi}{4}}/\mathbb{Z}_p) \leq \frac{2\pi^2 \cdot \lambda_p}{p}.$$

The first inequality follows from the Monotonicity Formula, Proposition 4.1 in the Appendix, applied to $T_{\frac{\pi}{4}} \subset (L(p,q), \lambda_p^2 g_{\mathbb{S}^3}, x)$. This finishes the proof of the lemma. □

Let's use the notation $\mathbb{Z}_p^q$ to represent the group $\mathbb{Z}_p$ acting on $\mathbb{S}^3$ and its dependence on the parameter q. By using the toroidal coordinate system for $\mathbb{S}^3$,

$$\varphi_r : \mathbb{R}^2 \to T_r : \varphi_r(u,v) = (\cos(r)e^{2\pi i u}, \sin(r)e^{2\pi i v}) \in \mathbb{S}^3,$$

the action of $\mathbb{Z}_p^q$ on T_r corresponds to the following action on $\mathbb{R}^2$:

$$(u,v) \longmapsto (u + \frac{kq}{p}, v + \frac{k}{p}).$$

In these coordinates , the $\mathbb{Z}_p^q$ orbit at the point $(z_0, w_0) = \varphi(\mathbb{Z} \times \mathbb{Z}) \in T_r$ is given by:

$$\text{Orbit}_{p,q}(z_0, w_0) = \{(m,n) + k(\frac{1}{p}, \frac{q}{p}) : m, n, k \in \mathbb{Z}\}. \tag{3.2}$$

Lemma 3.3. *Given a sequence $\{L(p,q)\}_{p \in \mathbb{N}}$, there exist $b, m_0, n_0 \in \mathbb{Z}$ and a subsequence $\{L(p_l, q_l)\}_{l \in \mathbb{N}}$ such that one of the following holds:*

1. *For every $(z_0, w_0) \in T_r$, $\varphi_r(Orbit_{p_l,q_l}(z_0, w_0))$ is becoming dense on T_r as $l \to \infty$.*

(2) $\varphi_r(Orbit_{p_l,q_l}(z_0,w_0))$ *is contained in* b *integral curves of* $X(z,w) = (m_0\sqrt{-1}\,z, n_0\sqrt{-1}\,w) \in \mathcal{X}(\mathbb{S}^3)$.

Proof. To prove the lemma it is enough to consider $(z_0,w_0) \in T_{\frac{\pi}{4}}$. If there is a subsequence for which the diameter of $T_{\frac{\pi}{4}}/\mathbb{Z}_p^q$ is going to zero as $p \to \infty$, then $\varphi_{\frac{\pi}{4}}(\mathrm{Orbit}_{p,q}(z_0,w_0))$ is clearly becoming dense on $T_{\frac{\pi}{4}}$ and item 1 is proved.

Let's consider now the case where the diameter of $T_{\frac{\pi}{4}}$ in $L(p,q)$ is bounded away from zero. From the equivalence between extrinsic and intrinsic distance and by Lemma 3.3 we conclude that the Euclidean injectivity radius satisfies $inj_{(z_0,w_0)}T_{\frac{\pi}{4}}/\mathbb{Z}_p = O(\frac{1}{p})$. In particular, there exist $k_p, m_p, n_p \in \mathbb{Z}$ such that

$$0 < ||k_p(\frac{1}{p},\frac{q}{p}) - (m_p,n_p)||_{\mathbb{R}^2} \leq \frac{C}{p}.$$

Therefore, there exists $(m_0,n_0) \in B_C(0) \cap \mathbb{Z}\times\mathbb{Z} \subset \mathbb{R}^2$ such that $(k_p - pm_p, k_pq - n_pp) = (m_0,n_0)$ infinitely often and $\frac{\sqrt{m_0^2+n_0^2}}{2p}$ is the Euclidean injectivity radius of $T_{\frac{\pi}{4}}/\mathbb{Z}_p$ at (z_0,w_0) for this subsequence. Hence, the sub-orbit generated by the translation $(u,v) \to (u,v) + k_p(\frac{1}{p},\frac{q}{p})$ is contained in the line $\mathbb{Z}\times\mathbb{Z} + \{t(m_0,n_0) : t \in \mathbb{R}\}$. It follows that the $\mathrm{Orbit}_{p,q}(z_0,w_0)$ is contained in a union of equidistant lines parallel to the one described above by homogeneity. Modulo $\mathbb{Z}\times\mathbb{Z}$ the number of such lines is finite, let's denote it by b_p. Modulo $\mathbb{Z}\times\mathbb{Z}$ there are $\frac{p}{b_p}$ points of $\mathrm{Orbit}_{p,q}(z_0,w_0)$ in each of these lines. Hence,

$$\frac{p}{b_p}(\frac{k_p}{p},\frac{k_pq}{p}) - \frac{p}{b_p}(m_p,n_p) = \frac{p}{b_p}(\frac{m_0}{p},\frac{n_0}{p}) \in \mathbb{Z}\times\mathbb{Z}.$$

Therefore, b_p divides m_0 and is independent of p. In other words, $\varphi_{\frac{\pi}{4}}(\mathrm{Orbit}_{p,q}(z_0,w_0))$ is contained in b integral curves of $X(z,w) = (m_0\sqrt{-1}z, n_0\sqrt{-1}w) \in \mathcal{X}(\mathbb{S}^3)$. □

Theorem 1.1. *There exists a positive integer* p_0 *such that for every* $p \geq p_0$ *and every* $q \geq 1$ *the isoperimetric surfaces in* $L(p,q)$ *are either geodesic spheres or tori of revolution about geodesics.*

Proof of Theorem 1.1. Arguing by contradiction, let's assume that there exists an infinite sequence of Lens spaces $L(p,q)$ containing an isoperimetric surface Σ_p of genus 2 or 3 for each p.

We use the Cheeger-Gromov convergence for the sequence of Lens spaces:

Cheeger-Gromov Convergence: A sequence (M_i,g_i,p_i) converges, in the sense of Cheeger-Gromov, to (M,g,p) if the following two conditions hold true:

(1) There exists an exhaustion of M by compacts Ω_i, i.e. $\Omega_i \subset \Omega_{n+1}$ and $\bigcup_i \Omega_i = M$.
(2) There exists a family of diffeomorphism onto their images, $\phi_i : \Omega_i \to M_i$, such that $\phi_i(p) = p_i$ and $\phi_i^* g_i \to g$ in the C^∞ topology.

We consider pointed manifolds $(L(p,q), p^2 g_{\mathbb{S}^3}, x_p)$ with base points x_p belonging to Σ_p. By the Cheeger-Gromov compactness theorem we have that $(L(p,q), p^2 g_{\mathbb{S}^3}, x_p) \to (M, \delta, x_\infty)$, where (M, δ) is a flat three manifold. The inclusion of Σ_p into M through the diffeomorphism ϕ_p is still denoted by Σ_p.

Lemma 3.4. *Let $(L(p,q), p^2 g_{\mathbb{S}^3}, x_p) \to (M, \delta, x_\infty)$ as above. There exists a constant $C > 0$ such that $|A_{\Sigma_p}|_{p^2 g_{\mathbb{S}^3}} \leq C$.*

Proof. Let $y_p \in \Sigma_p \subset L(p,q)$ be such that $|A_p|(y_p) = \max_{\Sigma_p} |A_p|^2$ and define $\lambda_p = \max_{\Sigma_p} |A_p|(y_p)$. Arguing by contradiction, let's assume that $\frac{\lambda_p}{p} \to \infty$. In local coordinates around y_p we consider the surface $\Sigma'_p = \lambda_p \Sigma_p$ on the Euclidean ball $B_{\lambda_p \frac{\pi}{10p}}(0)$ endowed with the rescaled metric $\lambda_p^2 g_{\mathbb{S}^3}$. Therefore, $(B_{\lambda_p \frac{\pi}{10p}}(0), \lambda_p^2 g_{\mathbb{S}^3}, y_p)$ converges to $(\mathbb{R}^3, \delta, 0)$ as $p \to \infty$. The surface Σ'_p now has the property that $\max_{\Sigma'_p} |A'_p(0)|^2 = 1$.

By the strong compactness for a sequence of isoperimetric surfaces with bounded second fundamental form, see Corollary 4.4 in the Appendix, there exists a subsequence converging to a properly embedded surface $\Sigma' \subset \mathbb{R}^3$, the convergence is in the sense of graphs and with multiplicity one. Moreover, Σ' is also stable, i.e.:

$$I_{\Sigma'}(f,f) \geq 0, \ \forall f \in C_0^\infty(\Sigma') \text{ satisfying } \int_{\Sigma'} f \, d_{\Sigma'} = 0.$$

If Σ' is compact, then it has to be a round sphere by Alexandrov's Theorem, which is a contradiction since strong convergence preserves topology. If Σ' is non-compact, then it has infinite area by the monotonicity formula: indeed, by Proposition 4.1 in the Appendix there exists a positive constant C such that

$$\frac{d}{dr}\left(\frac{e^{C\,r} \, |\Sigma' \cap B_r(x)|}{r^2} \right) \geq 0.$$

In particular, $|\Sigma' \cap B_r(x)| \geq \pi r^2$. As Σ' is properly embedded, it has infinite extrinsic diameter and the claim follows. Therefore, Σ' is totally geodesic by Da Silveira's Theorem 4.6 in the Appendix, which is a contradiction since $\max_{\Sigma'} |A| = 1$. □

Lemma 3.5. *Let $(L(p,q), p^2 g_{\mathbb{S}^3}, x_p) \to (M, \delta, x_\infty)$ as above, then M is a flat manifold of rank at most one.*

Proof. Below we denote $T_r/\mathbb{Z}_p$ by T_r. Let T_{r_p} be the Clifford torus through x_p enclosing a region Ω_{r_p}. Under the scaling by $\lambda_p = p^2$ we have that $|\Omega_{r_p}| = 2\pi^2 p^2 \sin^2(r_p)$. If $\lim_{p\to\infty} |\Omega_{r_p}| < \infty$, then $\lim_{p\to\infty} |T_{r_p}| = 2\pi^2 p \sin(2r_p) < \infty$. Moreover, the second fundamental form A_{r_p} of T_{r_p} satisfies $\lim_{p\to\infty} |A_{r_p}|^2 = \lim_{p\to\infty} \frac{1}{p^2}(\frac{\cos^2(r_p)}{\sin^2(r_p)} + \frac{\sin^2(r_p)}{\cos^2(r_p)}) < \infty$. The critical fiber $T_0 \subset \Omega_{r_p}$ is distant from x_p by $O(\frac{1}{p})$ since Ω_{r_p} is converging to a compact region in M. Instead of using base points x_p we choose new base points $y_p \in T_0$; it follows that $(L(p,q), p^2 g_{\mathbb{S}^3}, y_p) \to (N, \delta, y_\infty)$ and $\text{rank}(N) = \text{rank}(M)$. We claim that rank of N is at most one:

$$|B_{2R}(y_\infty)| = \lim_{p\to\infty} |B^p_{2R}(y_p)| \geq \lim_{p\to\infty} |\Omega_{\frac{R}{p}}| = \lim_{p\to\infty} 2\pi^2 p^2 \sin^2(\frac{R}{p}) = cR^2.$$

Let's assume now that $\lim_{p\to\infty} |\Omega_{r_p}| = \infty$, consequently $\lim_{p\to\infty} |T_{r_p}| = \infty$ and $\lim_{p\to\infty} |A_{r_p}|^2 = 0$. Recall the function $r = r(x)$, the distance from the Clifford torus through x to the critical fiber T_0 with respect to the round metric. The unit vector field ∂r is orthogonal to T_r for every r and it is well defined on $L(p,q) - \{T_0 \cup T_{\frac{\pi}{2}}\}$. Let $\gamma(r)$ be the geodesic whose velocity is ∂r and such that $\gamma(r_p) = x_p$. Consider $K_{r_p,R} = \{x \in T_r : d_{L(p,q)}(x, \gamma(r)) \leq \frac{R}{p} \text{ and } |r - r_p| \leq \frac{R}{p}\}$. By the triangle inequality $K_{r_p,R} \subset B_{2R}(x_p)$ under the metric $p^2 g_{\mathbb{S}^3}$. Applying the coarea formula for $f(r) = p\, r$, $|\nabla f|_{p^2 g_{\mathbb{S}^3}} = 1$, we obtain:

$$|K_{r_p,R}| = \int_{r_p - \frac{R}{p}}^{r_p + \frac{R}{p}} |B_R(\gamma(u)) \cap T_u| p\, du = |B_R(\gamma(u_0)) \cap T_{u_0}|_{p^2 g_{\mathbb{S}^3}} R \geq cR^2,$$

where $u_0 \in [r_p - \frac{R}{p}, r_p + \frac{R}{p}]$ is from the mean value theorem for integrals. The last inequality is justified as follows. Either the extrinsic diameter of T_{u_0} is going to infinity and T_{u_0} is converging with multiplicity to a flat surface or the extrinsic diameter of T_{u_0} is bounded. The former implies that $|B_R(\gamma(u_0)) \cap T_{u_0}|_{p^2 g_{\mathbb{S}^3}} \geq cR$. The latter implies that $B_R(\gamma(u_0)) \cap T_{u_0} = T_{u_0}$, which is a contradiction since $|T_{u_0}|_{p^2 g_{\mathbb{S}^3}} \to \infty$. We have conclude that $Vol(B_{2R}) \geq cR^2$ and rank of M is at most one. □

The following lemma gives a description of $I_{L(p,q)}$ for small volumes:

Lemma 3.6. *For p large enough there exist v_p and $\varepsilon_p > 0$ such that $I_{L(p,q)}$ is given by the profile of spheres on $(0, v_p]$ and by the profile of flat tori on $[v_p, v_p + \varepsilon_p)$. Moreover, if Σ_p is an isoperimetric surface such that $I_{L(p,q)}(v_p) = |\Sigma_p|$, then $g(\Sigma_p) = 0$ or 1.*

Proof. For each p we consider the first volume, v_p, for which there is transition on topology of isoperimetric surfaces from spheres to something else. If v_* is the volume for which the profile of geodesic spheres

intersect the profile of flat tori, then $v_p \leq v_*$. The value of v_* is computed by solving the following system of equations:

$$\frac{2\pi^2}{p}\sin^2(r) = 2\pi s - \pi\sin(2s) \quad \text{and} \quad \frac{2\pi^2}{p}\sin(2r) = 4\pi\sin^2(s).$$

The left hand sides (right hand sides) correspond to the enclosed volume and area of the Clifford torus T_r (geodesic spheres S_s of radius s), respectively. It follows that $s \leq \frac{\pi}{p}$; another way to see this is by recalling that the injectivity radius of $L(p,1)$ is $\frac{\pi}{p}$ at every point. Therefore, $I_{L(p,q)}(v_p) \leq O(\frac{1}{p^2})$.

Let Σ_p be an isoperimetric surface with the property that $I_{L(p,q)}(v_p) = |\Sigma_p|$. By Lemma 3.4 the sequence $\{\Sigma_p\}_{p\in\mathbb{N}}$ has bounded second fundamental form in $(L(p,q), p^2 g_{\mathbb{S}^3}, x_p)$; thus, it strongly converges to a properly embedded surface Σ of finite area in some orientable flat three manifold (M,δ) of rank at most one by Lemma 3.5. By the monotonicity formula, Proposition 4.1 in the Appendix, Σ is a closed surface. It follows that the pre-image $\widehat{\Sigma}$ of Σ in $\mathbb{R}^3$ is contained in a solid cylinder. Hence, $\widehat{\Sigma}$ is an union of round spheres by Alexandrov's Theorem or is a surface of revolution about the axis of the cylinder by Theorem 4.5 in the Appendix. Therefore, $g(\Sigma) = 0$ or 1. From the strong compactness for isoperimetric surfaces, see Corollary 4.4, we have that $g(\Sigma_p) = 0\,\text{or}\,1$, $v_p = v_*$ and the existence of the desired $\varepsilon_p > 0$. □

Claim. *Theorem 1.1 follows if we can show that the isoperimetric surfaces separating $L(p,q)$ in two regions of the same volume are tori.*

Proof. By the strong compactness for isoperimetric surfaces, Corollary 4.4, there exists $\widehat{v}$ such that if Σ is an isoperimetric surface enclosing volume $v \in [\widehat{v}, \frac{\pi^2}{p}]$, then Σ is a flat torus. It follows from Lemma 3.6 and for large p that the isoperimetric profile $I_{L(p,q)}$ is given by the area of geodesic spheres for volumes in $(0, v_p]$ and by the area of flat tori for volumes in $[v_p, \varepsilon_p] \cup [\widehat{v}, \frac{\pi^2}{p}]$. In other words, if $f(v)$ is the function defined by $f(v) = |T_{r(v)}|$, where $T_{r(v)}$ is the Clifford torus enclosing a volume equal to v, then $I_{L(p,q)}(v) = f(v)$ on $[v_p, \varepsilon_p] \cup [\widehat{v}, \frac{\pi^2}{p}]$. It follows that $\phi(v) = f(v) - I_{L(p,q)}(v)$ has a local maximum point at $t_* \in [\varepsilon_p, \widehat{v}]$.

The claim will follow by exploring the weak differential equation for $I_{L(p,q)}$. From (2.7) we have

$$f^2(v)f''(v) + \int_{T_{r(v)}} (2 + |A_{r(v)}|^2)\, d_{T_{r(v)}} = 0.$$

Let Σ be an isoperimetric surface such that $I_{L(p,q)}(t_*) = |\Sigma|$. If Σ_v is the unit normal variation of Σ parametrized by the enclosed volume, then

we define $h(v) = Area(\Sigma_v)$. Since $h \geq I_{L(p,q)}$, we have that $\phi_1 = f - h$ has also a local maximum point at t_*. Hence, $\phi_1'(t_*) = 0$, i.e., $H_r = H$, and $\phi_1''(t_*) \leq 0$. Applying equation (2.7) for h together with the Gauss equation and the Gauss-Bonnet theorem we obtain:

$$\begin{aligned}
\phi_1''(t_*) \leq 0 \quad &\Rightarrow \quad \frac{1}{h^2}\int_\Sigma (2+|A|^2)\, d_\Sigma \leq \frac{1}{f^2}\int_{T_{r(v)}} (2+|A_{r(v)}|^2)\, d_{T_{r(v)}} \\
&\Rightarrow \quad 4(1+H^2)\, h + 8\pi\,(g-1) \;\leq\; \left(4(1+H_r^2)\, f\right)\frac{h^2}{f^2} \\
&\Rightarrow \quad 1 + H^2 + \frac{2\pi(g-1)}{h} \leq (1+H_r^2)\,\frac{h}{f} \leq 1 + H^2.
\end{aligned}$$

Therefore, $g(\Sigma) = 1$ and $I_{L(p,q)} = f$ in $[v_p, \frac{\pi^2}{p}]$. □

By Lemma 3.3, the proof of Theorem 1.1 reduces to investigating the Cases I and II below.

Case I: There is a subsequence whose $\mathbb{Z}_p^q$ orbit of a point is contained in a finite number (independent of p) integral curves of a vector field $X(z,w) = (m_0\sqrt{-1}\, z, n_0\sqrt{-1}\, w) \in \mathcal{X}(\mathbb{S}^3)$.

We claim that the injectivity radius of $L(p,q)$ at every point is $O(\frac{1}{p})$. Indeed, let $\mathrm{Orbit}_{p,q}(z_0,w_0)$ be the orbit of $(z_0,w_0) \in \mathbb{S}^3$ with respect to $\mathbb{Z}_p^q$. As before, $\frac{p}{b}$ points of $\mathrm{Orbit}_{p,q}(z_0,w_0)$ lie on the curve $\beta(t) = \psi(z_0,w_0,t)$. Here, ψ is the one parameter family of diffeomorphisms associated to X:

$$\psi : \mathbb{S}^3 \times \mathbb{R} \to \mathbb{S}^3 : \psi(z,w,t) = (e^{m_0 it} z, e^{n_0 it} w).$$

When ordered according the orientation of $\beta(t)$, those points determine a piecewise closed geodesic $\gamma_p(t)$ with $\gamma_p(0) = (z_0,w_0)$. As $p \to \infty$, $\gamma_p(t)$ converges to $\beta(t)$. The claim now follows from :

$$\lim_{p\to\infty} \frac{p}{b} inj_{(z_0,w_0)} L(p,q) \leq \lim_{p\to\infty} \mathcal{H}^1(\gamma_p) = 2\pi\sqrt{m_0^2|z_0|^2 + n_0^2|w_0|^2}. \tag{3.3}$$

By the Cheeger-Gromov Compactness Theorem we obtain, up to subsequence, the following convergence:

$$(L(p,q),\, p^2\cdot g_{\mathbb{S}^3},\, x_p) \xrightarrow{C-G} (M,\, \delta,\, x_\infty), \tag{3.4}$$

where $(M,\, \delta,\, x_\infty)$ is a flat three manifold of rank one by Lemma 3.5. By (3.3) we conclude that $M = \mathbb{S}^1(r_0) \times \mathbb{R}^2 = \mathbb{R}^3/\mathbb{Z}$.

The curves $t \to \beta(t) = \psi(x,t)$ represent integral curves of X through $x \in L(p,q)$; they have bounded geodesic curvature and $\mathcal{H}^1(\beta) = O(\frac{1}{p})$. Hence, the integral curves of X converge to closed geodesics in M under (3.4). As sets, they coincide with the standard fibers of $M = \mathbb{S}^1 \times \mathbb{R}^2$.

By the Poincare-Hopf index theorem there exists a zero for the vector field $\frac{X^\top}{|X|} \in \mathcal{X}(\Sigma_p)$ since $g(\Sigma_p) \geq 2$. Hence, we can choose the base points x_p to satisfy $g_{\mathbb{S}^3}(\frac{X}{|X|}(x_p), N(x_p)) = \pm 1$, here N is the unit normal vector of Σ_p.

Lemma 3.7. *There exists a properly embedded surface $\Sigma_\infty \subset M$ such that $(\Sigma_p, x_p) \to (\Sigma_\infty, x_\infty)$ with multiplicity one. Moreover, Σ_∞ is totally geodesic and perpendicular to the standard fibers of $\mathbb{S}^1 \times \mathbb{R}^2$.*

Proof. By Lemma 3.4 the sequence of isoperimetric surfaces $(\Sigma_p, x_p) \subset (L(p,q), p^2 g_{\mathbb{S}^3}, x_p)$ has uniformly bounded second fundamental form. Applying the strong compactness theorem for isoperimetric surfaces, Corollary 4.4, we conclude that Σ_p converges smoothly and with multiplicity one to a properly embedded stable CMC surface $\Sigma_\infty \subset \mathbb{S}^1 \times \mathbb{R}^2$. If $\liminf_{p\to\infty} |\Sigma_p|_{p^2 g_{\mathbb{S}^3}} < \infty$, then the monotonicity formula, Proposition 4.1, implies that the extrinsic diameter of Σ_p and Σ_∞ are bounded. This is impossible since the sequence Σ_p separates $L(p,q)$ in two regions of the same volume that goes to infinity as $p \to \infty$. Therefore, Σ_∞ is a complete properly embedded stable CMC surface in $\mathbb{S}^1 \times \mathbb{R}^2$ with infinite area. Applying Da Silveira's Theorem 4.6 once more, we obtain that Σ_∞ is totally geodesic. As $g_{\mathbb{S}^3}(\frac{X}{|X|}(x_p), N(x_p)) = \pm 1$, we conclude that Σ_∞ is orthogonal to the standard fibers of $\mathbb{S}^1 \times \mathbb{R}^2$. □

We claim that Σ_∞ separates $\mathbb{S}^1 \times \mathbb{R}^2$. If it does not separate, then there exists a loop γ intersecting Σ_∞ at a single point. As $\Sigma_p \to \Sigma_\infty$ with multiplicity one, the same conclusion holds for Σ_p, which contradicts the fact that Σ_p separates $L(p,q)$. Therefore, there is $k \geq 1$ such that $\Sigma_\infty = \partial\Omega_\infty = \bigcup_{i=1}^{2k} \sigma_i$, where σ_i is a flat plane for each i.

Claim. *This configuration cannot be a limit of isoperimetric surfaces.*

We regard $\mathbb{S}^1 \times \mathbb{R}^2$ as a slab in $\mathbb{R}^3$ with height 2π. Now, we construct a deformation of Σ_∞ which decreases its area as follows. First, we cut off the k solid cylinders obtained from the intersection of Ω_∞ with a vertical solid torus of radius R. To balance the enclosed volume we add a vertical solid torus of radius r, see Figure 1 for the case $k = 1$. If a_i is the distance between σ_{2i} and σ_{2i-1}, then the radius r is given by:

$$\sum_{i=1}^{k} \pi\, R^2 a_i = \pi\, r^2 2\pi \Rightarrow r = R\sqrt{\frac{\sum_i a_i}{2\pi}}.$$

The boundary of this new region is denoted by $\widetilde{\Sigma}_\infty$ and

$$Area(\widetilde{\Sigma}_\infty \cap K) = Area(\Sigma_\infty \cap K) - 2k\pi\, R^2 + 2\pi\, R \cdot \sum_i a_i + 2\pi\, r \cdot 2\pi.$$

If R is large enough, then $\widetilde{\Sigma}_\infty$ has less area than Σ_∞. This is impossible



FIGURE 1. Compact support deformation of Σ_∞.

since the strong multiplicity one convergence allow us to carry out this deformation of Σ_∞ to Σ_p which contradicts the fact that Σ_p is an isoperimetric surface.

Case II: There is a subsequence $p \to \infty$ where the $\mathbb{Z}_p^q$ orbit of a point is becoming dense on the Clifford torus containing such point.

We use geometric measure theory methods to analyse the pre-image sequence $\{\Sigma'_p\}_{p\in\mathbb{N}} \subset \mathbb{S}^3$.

It is proved in [R2] that if $\Sigma = \partial\Omega$ has positive mean curvature, then

$$3\,|\Omega| \;\leq\; \int_\Sigma \frac{1}{H}\, d_\Sigma.$$

Applying this formula to the sequence of isoperimetric surfaces, we conclude that the mean curvature of Σ_p satisfy $H_p \leq \frac{|\Sigma_p|}{3|\Omega_p|} \leq \frac{2}{3}$. We have used that Σ_p separates $L(p,q)$ into regions of the same volume.

As $\{\Sigma_p\}_{p\in\mathbb{S}^3}$ has area and mean curvature bounded, we apply *Allard's compactness theorem*, Theorem 42.7 and Remark 42.8 in [S], to obtain an integral varifold $0 \neq \mathcal{V}^2 \subset \mathbb{S}^3$ that, up to subsequence, is $\mathcal{V}^2 = \lim_{p\to\infty} \Sigma'_p$. Recall that $\mathbb{S}^1\times\mathbb{S}^1$ acts on $\mathbb{S}^3$ via $(z,w) \to (\alpha_1,\alpha_2)(z,w) := (\alpha_1\, z, \alpha_2\, w)$. We claim that $\mathcal{V}^2$ is $\mathbb{S}^1 \times \mathbb{S}^1$ invariant: indeed, if $(z,w) \in \mathrm{supp}(\mathcal{V}^2)$ and $(\alpha_1,\alpha_2) \in \mathbb{S}^1\times\mathbb{S}^1$, then for each p there is $l_p \in \mathbb{Z}$ such that $\lim_{p\to\infty}(e^{\frac{2\pi i l_p q}{p}} z, e^{\frac{2\pi i l_p}{p}} w) = (\alpha_1 z, \alpha_2 w)$. On the other hand, as $(z,w) \in \mathcal{V}^2$, there is $(z_p, w_p) \in \Sigma'_p$ such that $(e^{\frac{2\pi i l_p q}{p}} z_p, e^{\frac{2\pi i l_p}{p}} w_p) \in \Sigma'_p$ and

$\lim_{p\to\infty}(z_p, w_p) = (z, w)$. It follows that $\lim_{p\to\infty}(e^{\frac{2\pi i l p q}{p}} z_p, e^{\frac{2\pi i l p}{p}} w_p) = (\alpha_1 z, \alpha_2 w)$ and $(\alpha_1, \alpha_2)(z, w) \in \mathrm{supp}(\mathcal{V}^2)$. In particular, $supp\, \mathcal{V}^2 = \bigcup_{j=1}^{k} T_{r_j}$ where T_{r_j} is a Clifford torus. The monotonicity formula implies that the convergence $\Sigma'_p \to \mathcal{V}^2$ is also in Hausdorff distance; hence, we have that $\Sigma'_p = \cup_{j=1}^{k} \Sigma'^j_p$ and $\mathrm{supp}\,(\lim_{p\to\infty} \Sigma'^j_p) = T_{r_j}$. Since Σ'_p is $\mathbb{Z}^q_p$ invariant there exists $\theta^{j_1 j_2}_p \in \mathbb{Z}^q_p \subset \mathbb{S}^1 \times \mathbb{S}^1$ for which $\theta^{j_1 j_2}_p(\Sigma'^{j_2}_i) = \Sigma'^{j_1}_p$. By taking the limit we obtain $(\alpha_1, \alpha_2)(T_{r_{j_2}}) = T_{r_{j_1}}$ for some $(\alpha_1, \alpha_2) \in \mathbb{S}^1 \times \mathbb{S}^1$. As this is impossible, we conclude that $k = 1$ and all Σ'_p are connected for p large.

Now we consider $\{\Sigma'_p\}_{p=1} \subset \mathbf{I}_2(\mathbb{S}^3, \mathbb{Z})$, the space of 2-dimensional integral *currents* on $\mathbb{S}^3$. Each $\Sigma'_p = \partial\Omega'_p$ and $\Omega'_p \in \mathbf{I}_3(\mathbb{S}^3, \mathbb{Z})$. As Ω'_p is a region of finite perimeter ($\mathcal{X}_{\Omega'_p}$ is BV function with uniform bounded variation), then $\Omega'_i \to \Omega'$ and $\Sigma'_p \to \partial\Omega'$ as currents, Ω' is an open set of finite perimeter, see Theorem 6.3 and proof of Theorem 37.2 in [S]. Since $|\Omega_p| = \pi^2$ we conclude that $|\Omega'| = \pi^2$. Applying the *Constancy theorem*, Theorem 26.27 in [S], we conclude that Ω' is the handlebody bounded by the Clifford torus T_{r_1}, and consequently $r_1 = \frac{\pi}{4}$.

We proved that $\mathcal{V}^2 = m\, T_{\frac{\pi}{4}}$ for some positive integer $m \in \mathbb{N}$. Since Σ_p is isoperimetric, it follows that $m = 1$. Indeed,

$$m\, |T_{\frac{\pi}{4}}| = |\mathcal{V}^2| = \lim_{p\to\infty} |\Sigma'_p| \leq |T_{\frac{\pi}{4}}|.$$

As $T_{\frac{\pi}{4}}$ is smooth, we have for $r > 0$ sufficiently small that the density $\theta(T_{\frac{\pi}{4}}, r, x) \leq 1 + \frac{\epsilon}{2}$, where $\epsilon > 0$ is from Theorem 4.2 in the Appendix. On the other hand, as Σ'_p is converging to $T_{\frac{\pi}{4}}$ with multiplicity one, then $\theta(\Sigma'_p, x, r) \leq 1 + \epsilon$ for p large enough. Now we invoke the smooth version of *Allard's Regularity Theorem*, Theorem 4.2, to concluded that the convergence $\Sigma'_p \to T_{\frac{\pi}{4}}$ is strong, i.e., graphical with multiplicity one. As strong convergence preserves topology, we conclude that $g(\Sigma_p) = 1$. This completes the proof of Theorem 1.1. □

3.3. **Berger Spheres.** Let g_0 be the round metric on $\mathbb{S}^3$ and J the vector field on $\mathbb{S}^3$ defined as $J(z, w) = (\sqrt{-1}\, z, \sqrt{-1}\, w)$. Recall that J is tangent to the fibers of the Hopf fibration $h : \mathbb{S}^3 \to \mathbb{S}^2(\frac{1}{2})$.

The *Berger metrics* are Riemannian metrics g_ε on $\mathbb{S}^3$ defined as:

$$g_\varepsilon(X, Y) = g_0(X, Y) + (\varepsilon^2 - 1) g_0(X, J) g_0(Y, J), \;\; \varepsilon > 0.$$

The Riemannian manifolds $(\mathbb{S}^3, g_\varepsilon)$ are called the *Berger spheres*, they are denoted by $\mathbb{S}^3_\varepsilon$. Geometrically, the metric g_ε shrinks the Hopf fibers to have length $2\pi\,\varepsilon$.

The Berger metrics are also homogeneous and their group of isometries has dimension four. It follows from the work of Abresch and

Rosemberg [AR] that every constant mean curvature surface in $\mathbb{S}^3_\varepsilon$ admits a holomorphic quadratic differential. In particular, every CMC sphere in $\mathbb{S}^3_\varepsilon$ is rotationally invariant.

A precise study of closed orientable surfaces with constant mean curvature on the Berger spheres is given in [TU]. It is proved there the existence of $\varepsilon_1 > 0$ with the following property: if $\varepsilon \in [\varepsilon_1, 1]$, then every stable constant mean curvature surface in $\mathbb{S}^3_\varepsilon$ has genus zero or one. Moreover, if $\varepsilon^2 \in [\frac{1}{3}, 1]$, then these stable CMC surfaces are totally umbilical spheres or the minimal Clifford torus, the latter only occurring when $\varepsilon^2 = \frac{1}{3}$. In particular, rotationally invariant spheres are the only solutions of the isoperimetric problem in $\mathbb{S}^3_\varepsilon$ for $\varepsilon^2 \in [\frac{1}{3}, 1]$.

Theorem 3.8. *There exists $\varepsilon_0 > 0$ such that for every $\varepsilon < \varepsilon_0$ the isoperimetric surfaces in the Berger spheres $\mathbb{S}^3_\varepsilon$ are either rotationally invariant spheres or tori.*

Proof. Let's assume that for every ε there exists an isoperimetric surface Σ_ε with $g(\Sigma_\varepsilon) = 2$ or 3.

We rescale the metric g_ε of $\mathbb{S}^3_\varepsilon$ by the factor $\lambda_\varepsilon = \frac{1}{\varepsilon^2}$. The Hopf fibers have constant length equal to 2π under the new metric $\lambda_\varepsilon g_\varepsilon$. It follows that the injective radius of $\mathbb{S}^3_\varepsilon$ at a point p is equal to $inj_p \mathbb{S}^3_\varepsilon = \pi$ for every $p \in \mathbb{S}^3_\varepsilon$.

Since h is a local trivial fibration, we have that for each $p_\varepsilon \in \mathbb{S}^3_\varepsilon$ there exist a neighbourhood V of $h(p_\varepsilon)$ and a diffeomorphism $\phi_\varepsilon : V \times \mathbb{S}^1 \to h^{-1}(V)$ such that $h \circ \phi_\varepsilon = \pi_1$, where $\pi_1 : V \times \mathbb{S}^1 \to V$ given by $\pi(x, y) = x$. Moreover, $\phi^*_\varepsilon(\frac{1}{\varepsilon^2} g_\varepsilon) \to \delta$ in the C^∞ topology. Therefore, in the sense of Cheeger-Gromov we have:

$$\big(\mathbb{S}^3_\varepsilon, \frac{1}{\varepsilon^2}\, g_\varepsilon, p_\varepsilon\big) \to \big(\mathbb{S}^1 \times \mathbb{R}^2, \delta, 0\big).$$

We pick the points $p_\varepsilon \in \Sigma_\varepsilon$ with the property that $g_\varepsilon(J, N_\varepsilon)(p_\varepsilon) = \pm\varepsilon$, this means J and N_ε are parallel at p_ε. These points exist by the Poincaré-Hopf index theorem. By Lemma 3.4 the inclusion of Σ_ε in $\big(\mathbb{S}^1 \times \mathbb{R}^2, \phi^*_\varepsilon(\varepsilon^{-2} g_\varepsilon)\big)$ has the following property:

$$\textit{There exists } C > 0 \textit{ such that } \sup_{\Sigma_\varepsilon} |A_\varepsilon| \leq C \text{ for every } \varepsilon.$$

By the strong compactness theorem for isoperimetric surfaces, Corollary 4.4 in the Appendix, we can extract a subsequence, $\{\Sigma_{\varepsilon_n}\}$, which converges with multiplicity one to a properly embedded surface $\Sigma_\infty \subset (\mathbb{S}^1 \times \mathbb{R}^2, \delta)$.

If $Area(\Sigma_\infty) < \infty$, then the monotonicity formula, Proposition 4.1, implies that Σ_∞ is compact. We apply Theorem 4.5 to conclude that Σ_∞ is either a round sphere or torus. This is impossible since we have

strong convergence and $g(\Sigma_\varepsilon) = 2\, or\, 3$. Therefore, Σ_∞ is a complete non-compact surface with infinite area. Moreover, Σ_∞ is also a stable CMC surface in $\mathbb{S}^1 \times \mathbb{R}^2$:

$$I_{\Sigma_\infty}(f,f) \geq 0, \;\; \forall f \in C_0^\infty(\Sigma_\infty) \text{ such that } \int_{\Sigma_\infty} f \, d_{\Sigma_\infty} = 0.$$

It follows from Theorem 4.6 that Σ_∞ is totally geodesic. By the choice of p_ε we conclude that Σ_∞ is orthogonal to the $\mathbb{S}^1$ fibers of $\mathbb{S}^1 \times \mathbb{R}^2$. Since Σ_∞ separates $\mathbb{S}^1 \times \mathbb{R}^2$, we also conclude that Σ_∞ is an union of at least two totally geodesic planes. As shown in the proof of Theorem 1.1, this configuration cannot be a limit of isoperimetric surfaces. □

4. Appendix

In this Appendix we collect some background results for surfaces with constant mean curvature in 3-manifolds.

Proposition 4.1. *Let M^3 be a three manifold with bounded curvature, $|K_M| \leq k$, and with positive lower bound on the injectivity radius $inj(M) \geq i_0$. If $\Sigma \subset M^3$ is a smooth surface with mean curvature which satisfies $|H| \leq H_0$, then there exists a positive constant $C = C(H_0, k)$ such that*

$$\frac{d}{dr}\left(\frac{e^{C\,r}\, Area(\Sigma \cap B_r(p))}{r^2}\right) \geq 0, \tag{4.1}$$

for every $p \in \Sigma$ and $r \leq \min\{i_0, \frac{1}{\sqrt{k}}\}$.

Proof. See Chapter 7 in [CM]. □

Let Σ be a CMC surface in a closed manifold M^3. The density of Σ at x is given by

$$\theta(\Sigma, x, r) = \frac{\text{Area}(\Sigma \cap B_r(x))}{\pi\, r^2}.$$

Theorem 4.2 (Allard's Regularity Theorem)**.** *Let M^3 be a closed manifold and $\rho > 0$. There exist $\epsilon = \epsilon(M, \rho) > 0$ and $C = C(M, \rho)$ with the following property: if $\Sigma \subset M$ is a smooth embedded CMC surface satisfying*

$$\theta(\Sigma, x, r) \leq 1 + \epsilon$$

for every $x \in M$ and $r < \rho$, then its second fundamental form is uniformly bounded, i.e., $|A_\Sigma| \leq C$.

Proof. See Theorem 1.1 in [W]. □

Let $\{\Sigma_n\}_{n\in\mathbb{N}}$ be a sequence of surfaces in a manifold M. We say that Σ_n converge to Σ in the sense of graphs if near any point $p \in \Sigma$ and for large n the surface Σ_n is locally a graph over an open set of $T_p\Sigma$ and these graphs converge smoothly to the graph of Σ. In addition, we say that $\{\Sigma_n\}$ satisfy *local area bounds* if there exist $r > 0$ and $C > 0$ such that $|\Sigma_n \cap B_r(x)| \leq C$ for every $x \in M$.

A hypersurface Σ is said to be *weakly embedded* if it admits only tangential self intersections.

Proposition 4.3. *Let $\{\Sigma_n\} \subset (M, g_n)$ be a sequence of embedded surfaces with constant mean curvature satisfying local area bounds and such that $\sup_{\Sigma_n} |A_n| \leq C$. Let's assume that g_n converges to a metric δ in the C^∞ topology. If $\{\Sigma_n\}_{n=1}$ has an accumulation point, then we can extract a subsequence that converges to a properly weakly embedded CMC surface Σ in (M, δ).*

Sketch of the Proof. Let's first recall the constant mean curvature equation for graphs. If Σ' is a surface with constant mean curvature H in (M, g), then Σ' can be written locally as a graph over a neighbourhood $U_p \subset T_p\Sigma'$:

$$\Sigma' = Graph(\, u\,) = \{(x_1, x_2, u(x_1, x_2) : x_1, x_2 \in U_p)\}.$$

In coordinates $g_{ij} := g(e_i, e_j)$ where $\{e_1, e_2, e_3\}$ is the coordinate base associated to (x_1, x_2, x_3). Let $\{E_1, E_2\}$ be the coordinate base for Σ_n, i.e., $E_i = e_i + u_{x_i}e_3 = T_i^l e_l$. The induced metric h is expressed by $h_{ij} = h(E_i, E_j)$. A simple computation gives:

$$g(N, e_i) = \frac{-u_{x_i}}{\sqrt{1 + g^{ij}u_{x_i}u_{x_j}}} \text{ and } g(N, e_3) = \frac{1}{\sqrt{1 + g^{ij}u_{x_i}u_{x_j}}}.$$

We also have

$$\begin{aligned} \nabla_{E_i}E_j &= T_i^l \nabla_{e_l} T_j^m e_m = T_i^l T_j^m \nabla_{e_l} e_m + E_i(T_j^m)e_m \\ &= T_i^l T_j^m \Gamma_{ml}^k e_k + u_{x_i x_j} e_3, \end{aligned}$$

where Γ_{ml}^k are the Christoffel symbols of g. Therefore,

$$g(N, \nabla_{E_i}E_j) = -\frac{\sum_{k=1}^2 T_i^l T_j^m \Gamma_{ml}^k u_{x_k}}{\sqrt{1 + g^{ij}u_{x_i}u_{x_j}}} + \frac{\left(u_{x_i x_j} + T_i^l T_j^m \Gamma_{ml}^3\right)}{\sqrt{1 + g^{ij}u_{x_i}u_{x_j}}}.$$

Finally, the mean curvature equation is written as:

$$H\sqrt{1 + g^{ij}u_{x_i}u_{x_j}} = h^{ij}u_{x_i x_j} + h^{ij}T_i^l T_j^m \Gamma_{ml}^3 - \sum_{k=1}^2 h^{ij}T_i^l T_j^m u_{x_k}\Gamma_{ml}^k. \tag{4.2}$$

Since $h_{ij} = T_i^l T_j^m g_{ml}$ and $T_i^3 = u_{x_i}$, then $|h^{ij}| \leq C_1(g^{ij}, u_{x_i}, u_{x_j})$. The equation (4.2) is uniformly elliptic as long as $|\nabla u|, |\nabla^2 u| < \widetilde{C}$.

Let $p \in M$ be an accumulation point for the sequence $\{\Sigma_n\}$. By the upper bound on the second fundamental form, $\sup_{\Sigma_n} |A_n| < C$, there exists $r_0 = r_0(C)$ such that for every $q \in B_{r_0}(p) \cap \Sigma_n$ we have that $\Sigma_n \cap B_{r_0}(q)$ is locally a graph u_n over a neighbourhood $U_q \subset T_q\Sigma_n$. Moreover, there exists $C_2 = C_2(C) > 0$ for which $\max\{\nabla u_n, \nabla^2 u_n\} \leq C_2$. As $|g_n - \delta|_{C^{2,\alpha}} \to 0$, the Schauder estimates for solutions of elliptic equations, see [GT], imply that u_n have $C^{2,\alpha}$ estimates on $B_{r_0/2}(q)$, i.e., $|u_n|_{C^{2,\alpha}} \leq C_3(|H_n| + |u_n|)$. Therefore, $u_n, \nabla u_n, \nabla^2 u_n$ are uniformly bounded and equicontinuous.

As $\{\Sigma_n\}_{n\in\mathbb{N}}$ satisfy local area bounds, then $|\Sigma_n \cap B(r_0/2)(y)| \leq C_4$. On the other hand, the monotonicity formula, Proposition 4.1, gives that $|\Sigma_n^j \cap B_{r_0}(y)| \geq C_5 r_0^2$, where Σ_n^j is a connected component of $\Sigma_n \cap B_r$. It follows that the number of components of $\Sigma_n \cap B_{r_0/2}(p)$ is finite and independent of n. By the Arzelà-Ascoli theorem we can extract a subsequence for which $\{u_n^j\}$ converges to u for every j. Moreover, u also satisfies the constant mean curvature equation (4.2). As the set of accumulation points of $\{\Sigma_n\}$ is compact in $B_R(p)$ we can cover this set by finite balls $B_{r_0}(p_k)$ with $k = 1, \ldots, N$. Repeating the arguments in each of these balls and applying a diagonal argument we obtain a properly immersed surface Σ on $B_R(p) \subset M$ with constant mean curvature H. Since the surfaces Σ_n are embedded, it follows that Σ does not cross itself though it may have tangential self-intersections. Therefore, Σ is properly weakly embedded in M. □

Corollary 4.4. *Let $(\Sigma_n, x_n) \subset (M_n, g_n, x_n)$ be a sequence of isoperimetric surfaces with $|A_n| \leq C$. Assume that (M_n, g_n, x_n) converges, in the sense of Cheeger-Gromov, to a three manifold (M, g, x). There exists a properly embedded surface $\Sigma \subset (M, g, x)$ such that $\Sigma_n \to \Sigma$ in the sense of graphs and the convergence is with multiplicity one.*

Sketch of the Proof. First we remark that $\{\Sigma_n\}$ satisfy local area bounds. Indeed, $\text{Area}(\Sigma_n \cap B_r(p)) \leq 2|\partial B_r(p)|$. By Proposition 4.3 we only need to rule out possibly multiplicities for the convergence $\Sigma_n \to \Sigma$ and points where Σ fails to be embedded.

If the multiplicity of the limit is bigger than two, then $\Sigma_n \cap B_r(p)$ has several components getting arbitrarily close. This allow us to do a local cut and past deformation, as shown in Figure 2, that preserves the enclosed volume. If δ is the Euclidean metric, then $\frac{1}{\widetilde{C}}\delta \leq g_n \leq \widetilde{C}\delta$ and $\frac{1}{C'}\text{Area}_\delta \leq \text{Area}_{g_n} \leq C'\text{Area}_\delta$. Thus, if $h \ll r$, then

$$Area_{g_n}(\Sigma_n') \leq Area_{g_n}(\Sigma_n) - C_1' r^2 + C_2' r h < \text{Area}_{g_n}(\Sigma_n).$$



FIGURE 2. Example of higher multiplicity

The deformation needed for the multiplicity two case is shown in Figure 3. The constraint on the enclosed volume implies that $\frac{4}{3}\pi R^3 \approx \pi r^2 h$. Hence, if $h \ll r$, then

$$Area(\Sigma'_n) \leq Area(\Sigma'_n) - C'_1 r^2 + C'_2 r\, h - C'_3 R^2 + C'_4 R^2 < Area(\Sigma'_n).$$

The construction to deal with points where Σ has tangential self intersections is similar to the multiplicity two case. The corollary now follows since these constructions contradict the fact that Σ_n is isoperimetric for every n. □



FIGURE 3. Example of multiplicity two

Theorem 4.5 (Korevaar-Kusner-Solomon)**.** *If Σ is a complete CMC surface properly embedded in a solid cylinder in $\mathbb{R}^3$, then Σ is rotationally symmetric with respect to a line parallel to the axis of the cylinder.*

Proof. See [KKS]. □

Theorem 4.6 (Da Silveira)**.** *Let (Σ, ds^2) be a complete orientable surface conformally equivalent to a compact Riemann surface punctured at a finite number of points. Let $L = \Delta + q$ be an operator on Σ satisfying*

$q \geq 0$ and $q \neq 0$. If Σ has infinite area, then there exists a piecewise smooth function f with compact support such that

$$-\int_\Sigma f\, L\, f\, d_\Sigma < 0 \quad \textit{and} \quad \int_\Sigma f\, d_\Sigma = 0.$$

Proof. See [DS]. □

References

[A] F. J. Almgren. *Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints*, Mem. AMS 165 (1976).

[AR] U. Abresch and H. Rosenberg. *Generalized Hopf differentials.* Mat. Contemp. 28 (2005), 1-28.

[B] H. Bray, *The Penrose inequality in general relativity and volume comparison theorems involving scalar curvature* (thesis). arXiv:0902.3241v1, (1998).

[BC] J.L. Barbosa and M. do Carmo. *Stability of hypersurfaces with constant mean curvature*, Math. Z., 185, 339-353, 1984.

[BCE] J.L. Barbosa, M. do Carmo, and J. Eschenburg. *Stability of hypersurfaces with constant mean curvature in Riemannian manifolds*, Math. Z., 197, 123-138, 1988.

[BM] H. Bray and F. Morgan. *An isoperimetric comparison theorem for Schwarzschild space and other manifolds*, Proc. Amer. Math. Soc. 130:5 (2002), 1467-1472.

[BP] C. Bavard and P. Pansu. *Sur le volume minimal de R^2*, Ann. Sci. Ecole Norm. Sup. (4) 19 (1986), 479-490.

[C] B.Y. Chen. *Geometry of submanifolds.* Pure and Applied Mathematics, No. 22. Marcel Dekker, Inc., New York, 1973. vii+298 pp.

[CM] T. Colding and W. Minicozzi. *A course in minimal surfaces.* Graduate Studies in Mathematics, 121. American Mathematical Society, Providence, RI, 2011. xii+313 pp.

[DS] A. Da Silveira. *Stability of complete noncompact surfaces with constant mean curvature.* Math. Ann. 277 (1987), no. 4, 629-638.

[GMT] E. Gonzalez, U. Massari and I. Tamanini. *On the regularity of boundaries of sets minimizing perimeter with a volume constraint*, Indiana Univ. Math. J. 32 (1983), 25-37.

[GT] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order.* Second edition. Fundamental Principles of Mathematical Sciences, 224. Springer-Verlag, Berlin, 1983. xiii+513 pp.

[HH] W. Hsiang and W. Hsiang. *On the uniqueness of isoperimetric solutions and imbedded soap bubbles in noncompact symmetric spaces*, I. Invent. Math. 98(1), (1989), 39-58.

[HPRR] L. Hauswirth, J. Pérez, P. Romon and A. Ros. *The periodic isoperimetric problem.* Trans. Amer. Math. Soc. 356 (2003), no. 5, 2025-2047.

[KKS] N.J. Korevaar, R. Kusner and B. Solomon. *The structure of complete embedded surfaces with constant mean curvature*, J. Differential Geom. 30 (1989), 465-503.

[LY] P. Li and S.T. Yau. *A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces*, Invent. Math., 69, 269-291, 1982.

[M] F. Morgan. *Regularity of isoperimetric hypersurfaces in Riemannian manifolds*, Trans. Amer. Math. Soc. 355(2003), Pages 5041-5052.

[MHH] F. Morgan, S. Howe and N. Harman. *Steiner and Schwarz symmetrization in warped products and fiber bundles with density*, Rev. Mat. Iberoamericana 27(2011), no.3, 909-918.

[MJ] F. Morgan and D. Johnson. *Some sharp isoperimetric theorems for Riemannian manifolds*. Indiana Univ. Math. J. 49 (2000), no. 3, 1017-1041.

[MN] F.C. Marques and A. Neves. *Min-max theory and the Willmore conjecture* Annals of Math. 179 2 (2014), 683-782.

[P] R. Pedrosa. *The isoperimetric problem in spherical cylinders*. Ann. Glob. Anal. Geom. 26(4), (2004), 333-354.

[PP] P. Petersen. *Riemannian geometry*. Second edition. Graduate Texts in Mathematics, 171. Springer, New York, 2006. xvi+401 pp.

[PR] R. Pedrosa and M. Ritoré. *Isoperimetric domains in the Riemannian product of a circle with a simply connected space form and applications to free boundary problems*. Indiana Univ. Math. J. 48(4), (1999), 1357-1394.

[R] A. Ros. *The Willmore conjecture in the real projective space*, Math. Res. Lett. 6 (1999), 487-493.

[R1] A. Ros. *The isoperimetric problem*, Proc. Clay Research Institution Summer School, 2001.

[R2] A. Ros. *Compact hypersurfaces with higher order mean curvatures*, Revista. Matematica. Iberoamericana. Vol 3, 1987, 447-453.

[RR] M. Ritoré and A. Ros. *Stable Constant Mean Curvature Tori and the Isoperimetric Problem in Three Space Forms*, Comment. Math. Helvet. 67 (1992), 293-305.

[RR1] M. Ritoré and A. Ros. *The spaces of index one minimal surfaces and stable constant mean curvature surfaces embedded in flat three manifolds,* Trans. Amer. Math. Soc. 348 (1996), 391-410.

[S] L. Simon. *Lectures on geometric measure theory*. Proceedings of the Centre for Mathematical Analysis, Australian National University 1983.

[SS] R. Schoen and L. Simon. *A new proof of the regularity theorem for rectifiable currents which minimize parametric elliptic functionals*. Indiana Univ. Math. J. 31 (1982), no. 3, 415-434.

[TU] F. Torralbo and F. Urbano. *Compact stable constant mean curvature surfaces in homogeneous 3-manifolds*. Indiana Univ. Math. J. 61 (2012), no. 3, 1129-1156.

[W] B. White. *A local regularity theorem for mean curvature flow*. Ann. of Math. (2) 161 (2005), no. 3, 1487-1519.

University College London UCL Union Building, 25 Gordon Street, London WC1E 6BT

E-mail address: c.viana@imperial.ac.uk